

SPECIFICATIONS

3Tac-Iβ

Active ingredient	3 live species of Trichoderma. 10mg (0,01 gr.) Clavulanic Ácid
Concentration and formulation	Trichoderma – no less than 25.000 ufc/gr., 30% of active Trichoderma per kilogram of weight and 70% of pulverized micro-nutrients per kilogram of dry product weight. 10mg (0,01 gr.) of Clavulanic Ácid
Controlled organisms	Gram Bacterias (+ and -), coniforms, Pseudomonas, Erwinia, Xantomonas, Canker and bacterial leaf streak. Secondary control of phytopathogenic fungi in primary stages
Action mechanism	Bactericide
Action on arthropods	None
Time to product activation	Rapid activation (6 hours from application)
Dose per application	2 kilograms per hectare in mild to moderate bacteriosis diluted in 600 liters of water. 3 or more kilograms per hectare in severe bacteriosis, diluted in 600 liters of water
Residual effect	20 days
Toxicity	Group IV. Products that are normally not harmful
Application equipment	Any application equipment
Application temperature	Range from 0° to 35° C
Restrictions	No entrance restrictions to orchards
Water and pH type	Can be used with any type of water and any pH level
Physical product	1 Kilogram of wetting powder
Storage	In a fresh and dry environment. Sealed container. No direct exposure to the sun.